

Izoconazol-M 600 mg

Pharmaceutical form: Vaginal suppository.

Composition: 1 suppository contains: active ingredient: isoconazole nitrate 600 mg.

Inactive ingredients: Polyethylene glycol 1500 and Polyethylene glycol 400.

Pharmacological group: antifungal agent.

Pharmacological properties:

Isoconazole is a synthetic derivative of imidazole. It has a local antifungal and antibacterial effect. It acts fungistatically. Active against dermatophytes, molds, yeasts, and yeast-like fungi, as well as gram-positive bacteria (staphylococci, micrococci, streptococci). Inhibits the synthesis of ergosterol and changes the composition of the microbial cell membrane.

Indications:

Fungal infections of the vagina, incl. mixed infections caused by gram-positive bacteria.

Contraindications:

Hypersensitivity to the active substance or to any of the excipients.

Posology and method of administration:

A vaginal suppository is inserted deep into the vagina. It is best to do this in the supine position, in the evening just before bedtime, 1 time/day once. For making easy to insert, fingerstalls should be used.

If necessary, you can repeat the course of treatment after 7-10 days.

Treatment should not be carried out during menstruation.

Undesirable effects:

When using the drug, local reactions may occur, such as irritation, burning, and itching. In rare cases, it is possible to develop allergic reactions at the site of the application of the drug.

Precautions and Warnings:

It is not recommended to douche during treatment, as well as within a week after treatment. To prevent re-infection, it is recommended to change and boil personal linen daily (towels, underwear - preferably made of cotton).

Simultaneous treatment of the sexual partner is recommended.

Pregnancy and Lactation:

The experience with the use of isoconazole-containing preparations during pregnancy does not indicate a teratogenic risk in humans.

Because of the only slight absorption (less than 5% of the dose administered) and the short duration of the therapy, damage to the baby from isoconazole nitrate transferred with the breast milk can be ruled out.

Package:

1 vaginal suppository in a strip with 2 fingerstalls, 1 strip and leaflet insert in a paperboard box.

Shelf life:

3 years.

Storage:

Store below 30 °C. Keep out of reach of children.

Legal status:

Prescription only.